

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 9/14, 9/00	A1	(11) International Publication Number: WO 93/13752 (43) International Publication Date: 22 July 1993 (22.07.93)
(21) International Application Number: PCT/US93/00373 (22) International Filing Date: 20 January 1993 (20.01.93) (30) Priority data: 07/823,218 21 January 1992 (21.01.92) US (71) Applicant: SRI INTERNATIONAL [US/US]; 333 Ravenswood Avenue, Menlo Park, CA 94025-3493 (US). (72) Inventors: PLATZ, Robert, M. ; 324 Valdez, Half Moon Bay, CA 94019 (US). IP, Anna ; 3649 South Court, Palo Alto, CA 94306 (US). WITHAM, Clyde, L. ; 1294 Birch Street, Montara, CA 94037 (US). (74) Agents: LANGE, Richard, P.; SRI International, 333 Ravenswood Avenue, Menlo Park, CA 94025-3493 (US) et al.		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: IMPROVED PROCESS FOR PREPARING MICRONIZED POLYPEPTIDE DRUGS (57) Abstract Solid particle aerosol formulations of polypeptide drugs are made by lyophilizing solutions of the drugs which contain milling stabilizers that inhibit degradation of the drug during subsequent milling. The lyophilized drug is milled in fluid energy mills that have been fitted with abrasion-resistant materials and which use pure nitrogen that has been filtered to eliminate particles of greater than 0.1 µm to transport the drug. The use of (a) milling stabilizers in the solution and (b) abrasion-resistant fluid energy mills that use pure filtered nitrogen in the milling step reduce insoluble contaminants and inactive fractions in the milled product.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

-1-

5 IMPROVED PROCESS FOR PREPARING
 MICRONIZED POLYPEPTIDE DRUGS

Description

10 Technical Field

 The present invention is in the field of
 pharmaceutical formulation. More particularly, it
 concerns improvements in a process for micronizing
 polypeptide drugs into powders that are suitable for
15 formulation as solid particle aerosols.

Background

 Inhalation therapy involves the administration
 of a drug in an aerosol form to the respiratory tract.
20 Two types of aerosols are employed: liquid particles and
 solid particles. The liquid aerosols are generated by
 nebulizing solutions of the drug. Solid particle
 aerosols are either in the form of a powder suspended in
 a propellant which is administered from a metered dose
25 inhaler or simply as a powder that is administered from a
 dry powder inhaler. In the case of polypeptide drugs,
 solid particle aerosols are typically made by
 lyophilizing the drug from solution and then milling or
 grinding the lyophilized drug to the desired particle
30 size distribution for pulmonary administration.

 Prior workers have recognized certain stability
 problems in making such drugs. First, prior workers have
 recognized that removal of residual water during the
 lyophilization in the presence of protective agents
35 provides a more stable lyophilized product. A variety of

-2-

excipients have been proposed as such cryoprotectants (Pikal, M.J., Biopharm. (1990) 3(8):18-27 and Franks, F., Cryoletters (1990) 11:93-110.

Prior workers have also recognized and
5 addressed problems associated with the "stability" of drug powders that are used in solid particle aerosols. These "stability" problems involve the particles being adsorbed or absorbed by materials in which they come into contact during their handling or the particles forming
10 agglomerates. For instance, PCT Application Publication No. WO90/09787 suggests using "extender" molecules to avoid particle adsorption, absorption, or agglomeration while metering the particles into unit dose containers. Among the suggested extenders are amino acids
15 (particularly DL-methionine), mono-, di-, and polysaccharides, and polypeptides.

To the best of applicants' knowledge, the art has not recognized or addressed the deleterious effects of the grinding or milling step of the powder production
20 on the solubility and biological activity of the product powder. In this regard, the present invention is directed to a means for reducing the amounts of insoluble material and biologically inactive material produced when a polypeptide drug is lyophilized and milled into a
25 powder for administration as a solid particle aerosol.

Disclosure of the Invention

The invention relates to improvements in a process for preparing a micronized polypeptide drug in
30 which a solution of the drug is lyophilized and the lyophilized drug is milled to a powder in a fluid energy mill using a pressurized feed gas. One improvement comprises using a purified inert feed gas which has been filtered to remove particles greater than about 0.1 μm
35 from the feed gas whereby water insoluble contaminants in

-3-

the powder are reduced. Another improvement applies to instances where the drug is labile under the milling conditions and comprises incorporating a milling stabilizer in the polypeptide drug whereby the production of biologically inactive material in the milling step is reduced.

Modes for Carrying Out the Invention

The present invention is useful for transforming polypeptide drugs into a powder form that is suitable for aerosol administration or for use in injectable suspensions. As used herein, the term "polypeptide" intends molecules composed of about 10 or more amino acids linked by peptide bonds. The polypeptides may include moieties, e.g., sugars other than amino acids, or be complexed with other moieties such as lipids. Examples of such polypeptides are insulin, interferon α , interferon β , interferon γ , interleukin 1-7, granulocyte colony stimulating factor (GCSF), macrophage granulocyte colony stimulating factor (MGCSF), epidermal growth factor (EGF), insulin-like growth factors 1 and 2 (IGF-1 and 2, transforming growth factor α (TGF- α), transforming growth factors β 1-3 (TGF- β 1, 2 and 3), tissue plasminogen activator (TPA), platelet-derived growth factor (PDGF), erythropoietin (EPO), human growth hormone (HGH), and other hormones, growth factors, and neuropeptides. These polypeptides may be prepared by isolating them from natural sources (body tissues or fluids), by synthesizing them by recombinant DNA technology, or in the case of relatively small molecules, by synthesizing them by solid-phase techniques with or without ligation.

The first step in the process for forming the polypeptides into micronized particles is lyophilization. If the polypeptide as isolated or synthesized is in

-4-

solution, no solvent need be added. If it is in a dry state, the polypeptide is dissolved in an appropriate solvent. In many instances, the polypeptide is water soluble. It will be appreciated that low solubility polypeptides simply require use of a greater volume of solvent. The polypeptide is lyophilized from a frozen solution. If the polypeptide is labile to the conditions to which it is subjected during milling (i.e., shear, pressure, temperature), a milling stabilizer is added to the solution. Such stabilizers include sugars, e.g., sorbitol, mannitol, sucrose, lactose, and trehalose, surfactants such as polyvinylpyrrolidone, and water-soluble proteins such as serum albumins (bovine, human, chicken, goat, etc.). The stabilizer will normally be added to the solution at concentrations of 0.5 to 50 mg/ml, preferably 1 to 20 mg/ml. Sorbitol in combination with other sugars is a preferred stabilizer. Alternatively, the stabilizer may be mixed homogeneously with the lyophilized cake. It will be appreciated that the optimum stabilizer or combination of stabilizers will vary from drug to drug and that the optimum stabilizer(s) and amount of stabilizer for a given polypeptide drug may need to be determined empirically.

The solution is lyophilized by freezing the solution and subjecting the frozen mass to a vacuum. Conventional lyophilization equipment may be used. Normally, the solution will be cooled to -10°C to -60°C, preferably -15°C to -45°C and subjected to a vacuum of 50 to 750 millitorr, preferably 100 to 250 millitorr for 50 to 120 hours.

The lyophilized polypeptide is then size reduced in a grinding mill, preferably a fluid energy mill also known as a jet mill. These mills employ high velocity gas streams to carry feed particles into contact with each other and/or the surfaces of a grinding

-5-

chamber. In one type the fluid energy is admitted via a multiplicity of high velocity gas streams at an angle around a portion or all of the periphery of a cylindrical milling-classifying chamber. In another type, fluid

5 streams convey the particles at high velocity into a chamber where two streams impact upon each other. In another type the fluid stream conveys the particles against a stationary impact surface or around a toroidal channel. In all types there is a high energy release and

10 a high order of turbulence which causes the particles to grind upon themselves and upon the mill surface and to be fractured. Most mills classify particles centrifugally using the energy of the fluid stream and have means for returning oversize particles to the grinding chamber for

15 further size reduction. Mills that employ the angle-jets are preferred. Examples of commercial embodiments of such mills are the Micronizer (Sturtevant Mill Corp.), the Jet Pulverizer (Jet Pulverizer Co.), the Reductionizer, the Micron-Master, the Trost M.N., and

20 the Jet-O-Mizer. Structurally these mills comprise a cylindrical grinding chamber wherein the lyophilized polypeptide drug is acted upon by a number of gas jets issuing through orifices spaced around the periphery of the chamber. The rotating gas discharges at the center

25 of the chamber carrying the fines with it, while the coarse particles are thrown toward the chamber wall where they are subjected to further reduction by impact from particles and gas in the incoming jets. The outlet from the grinding chamber normally leads directly into a

30 centrifugal product collector.

In order to reduce the amount of insoluble contaminants (i.e., materials that are insoluble in water at 37°C) present in the product powder, applicants have modified the conventional milling procedure in several

35 important respects. First, any parts of the mill (e.g.,

-6-

solder or weld joints) that may be subject to abrasion and degradation are replaced with materials that are more resistant to abrasion and degradation under the milling condition. Second, an inert feed gas (e.g., nitrogen) is used to fluidize the particles. In this regard, the term "inert" intends that the gas will not react with the lyophilized feed material under the milling conditions. Liquid nitrogen is a preferred source of inert gas because it is ultrapure and it helps cool the mill and the product. Third, the incoming feed gas is filtered prior to introduction into the mill to eliminate contaminants of less than about 0.1 μm particle size. Filtration may be accomplished by placing commercially available ULPA (ultrahigh efficiency) filters (e.g., Wafergard, Millipore, Inc.) in the incoming gas lines. Use of high purity gas lines is desirable to avoid introducing contaminants into the polypeptide.

As indicated above, applicants further employ milling stabilizers to reduce shear, pressure and/or temperature degradation of the polypeptide drug. These stabilizers may also protect the polypeptide from degradation during the lyophilization and/or during storage.

The particle size of the milled powder is primarily dependent upon the energy imparted by the gas to the particles which in turn is related to the pressure of the feed gas. This pressure will normally range between about 10 to 120 psig, more usually 80 to 110 psig. In general, lower pressures in this range (i.e., 10 to 30 psig) will produce larger particles (i.e., 4 to 15 μm in diameter) whereas higher pressures will produce smaller particles (i.e., 0.5 to 4 μm in diameter).

The desired particle size for a given polypeptide drug will depend upon the site in the respiratory tract for which the drug is targeted.

-7-

Generally, particles below about 4 μm in diameter (usually 0.5 to 4 μm) will deposit in the therapeutically effective lower pulmonary regions when administered as an aerosol into the respiratory tract whereas larger particles (4 to 10 μm) will lodge in the upper airways of the respiratory tract. It will be appreciated that the particle sizes recited herein refer to a mass distribution of particles defined by the mass median diameter (where 50% of the mass of the particles is above the stated size and 50% of the mass is below the stated size). Particle sizes may be determined by commercially available particle size analyzers.

By using the improved process of the invention, insoluble contaminants in the powder product are reduced to below about 5% by weight, more usually below about 2% by weight. These contaminants have been found to increase degradation of the active ingredient in the milled powder product. In the case of polypeptides that are otherwise degraded by jet milling, the invention reduces the extent of degradation such that the polypeptide drug component of the milled powder constitutes at least about 85% active drug, preferably at least about 95% active drug.

The milled powder may be formulated with a propellant for administration via metered dose inhaler devices. The propellant will normally be a hydrocarbon or halohydrocarbon such as tetrafluoroethane, trichlorofluoromethane, dichlorofluoromethane, and the like. Surfactants may be added to the formulation to facilitate and stabilize the suspension and dispersion of the powder in the propellant and to lubricate inhaler components. The drug powder will normally constitute about 0.1 to 10% by weight of the formulation, more usually 0.5 to 2.5% by weight.

35

-8-

The milled powder may be formulated with bulking agents or administered neat from powder inhaler devices. When bulking agents are used, they will normally constitute about 50% to 99.9% of the formulation.

The solid particle aerosol of the polypeptide drug will be administered in one or more unit doses to provide therapeutic levels of the drug. The amount of polypeptide administered and the dosing regimen will depend upon the particular drug being administered, the indication being treated, and the patient. Accordingly, it will be appreciated that the specific amount and regimen used in given instances may vary widely.

The following examples further illustrate the invention. These examples are not intended to limit the invention in any manner.

Example 1: Micronization of Human Growth Hormone

20

Lyophilization

Lyophilized recombinant human growth hormone was obtained from the manufacturer of same.

25

Milling

A Sturtevant Micron-Master mill (model 2") was used. Originally the feed gas to the mill was fed in at 100 psig. The insoluble fraction in the milled product was determined by U.V. spectrophotometer. In initial runs approximately 40% by weight of the milled product was insoluble.

The insoluble fraction was reduced to 20% by weight by (1) replacing the silver soldered joints in the mill with welded joints and (2) replacing the soldered copper gas lines with flexible stainless steel hose. The

-9-

insoluble fraction was further reduced to 5 to 15% by weight by (1) replacing the Sturtevant gas manifold with a manifold made of electropolished stainless steel tubing and VCR couplings, (2) installing a Balston particle filter (type BK tested for removing 0.1 μm particles at 99.99% efficiency), and (3) milling with 99.99% pure nitrogen. Further reduction of insoluble material was achieved by altering the powder-feed mechanism to prevent it from drawing contaminated room air and by supplying the gas from liquid nitrogen dewar containers. The insoluble fraction was further reduced to 2.5 to 7.5% by weight by replacing the gas manifold with a high purity gas manifold and ultra high purity filters (Wafergard, Millipore, Inc.).

15

Example 2: Micronization of IFN- β

Lyophilization

Aqueous solutions of IFN- β having the following compositions were prepared.

20

<u>Component</u>	<u>Solution A</u>	<u>Solution B</u>	<u>Solution C</u>
IFN/ β	15 $\mu\text{g/mL}$	15 $\mu\text{g/mL}$	65 $\mu\text{g/mL}$
Human Serum	1.5 mg/mL	15 mg/mL	1.445 mg/mL
25 Albumin			
NaCl	0.6 mg/mL	0.6 mg/mL	0.584 mg/mL
Sorbitol	-	4.5 mg/mL	-

The solutions were lyophilized in a freeze dryer (Model No. GT4, Leybold, Cologne, Germany)

30

35

-10-

Milling

The jet mill described in Example 1 that provided the best reduction of insolubles was used. The filter on the feed gas line was obtained from Millipore Corp. (Bedford, MA).

The mill was operated at 80-100 psig according to manufacturer's instructions to provide a powder having greater than 50% of the particles less than 3 μ m in diameter and 90% of the particles less than 5 μ m in diameter.

The lyophilized powder (before micronization) and the micronized powders were assayed for IFN- β activity using an enzyme immunoassay (EIA) (Yamazaki et al., J. Immunoassay (1989) 10:57). The results were as follows:

<u>Solution</u>	<u>Activity Before Micronization</u>	<u>Activity After Micronization</u>
A	107 \pm 17%	65 \pm 4.8%
B	110 \pm 25%	102 \pm 3.8%
C	10.4 \pm 2.5%	8.2 \pm 1.9%

As shown, the formulation that contained both human serum albumin and sorbitol withstood the milling without significant loss of activity.

Example 3: Micronization of GCSF

Aqueous formulations of GCSF at 1 mg/mL, glycine HCl at 1.1 mg/mL, and mannitol, lactose or sucrose at 10 mg/mL were made, lyophilized and milled. Pure nitrogen, milling pressure 100 psig, was used with a feed rate of the lyophilized material of 0.5 g/min. The milling apparatus was that used in Example 2 above. The milled powders contained insignificant amounts of

-11-

insoluble material but significant amounts of soluble aggregates.

In an attempt to eliminate soluble aggregates, an aqueous formulation of GCSF at 1.0 mg/mL, mannitol at 9.0 mg/mL and sorbitol at 1.0 mg/mL was made, lyophilized and milled as above. The addition of sorbitol reduced the level of soluble aggregates in the milled powder to $4\% \pm 1.87$. Samples of unmilled powder and milled powder of this formulation were reconstituted in glycine-buffered water (pH 3.5) and analyzed by size-exclusion HP chromatography. The resulting chromatographs indicated there was insignificant degradation of the powder by the milling.

Modifications of the above-described modes for carrying out the invention that are obvious to those of skill in the field of pharmaceutical formulation and related fields are intended to be within the scope of the following claims.

-12-

Claims

1. In a process for preparing a micronized polypeptide drug comprising lyophilizing a solution of the drug in a solvent and milling the lyophilized drug in a fluid energy mill using a pressurized gas, the improvement comprising using a pure inert gas filtered to remove particles greater than about 0.1 μm in diameter as the gas, whereby insoluble contaminants in the powder are reduced.

2. The process of claim 1 wherein the polypeptide drug is labile under the milling conditions and the lyophilized drug contains a milling stabilizer.

3. The process of claim 2 wherein the milling stabilizer comprises sorbitol, mannitol, sucrose, lactose, trehalose, serum albumin, polyvinylpyrrolidone, or mixtures thereof.

4. The process of claim 1 wherein the inert gas is nitrogen of greater than about 99% purity.

5. The process of claim 4 wherein the pressure of the nitrogen is in the range of 10 to 120 psig.

6. The process of claim 5 wherein the source of the nitrogen is liquid nitrogen.

7. The process of claim 1 wherein the particle size of the micronized polypeptide drug is in the range of about 0.5 to about 15 μm .

8. The process of claim 1 wherein the solvent is water.

-13-

9. The process of claim 1 wherein the parts of the mill that are contacted by the pressurized gas in which the polypeptide drug is being carried are not substantially abraded by said drug-carrying gas.

5

10. The process of claim 1 wherein the micronized polypeptide drug is suitable for administration as a solid particle aerosol.

10

11. In a process for preparing a micronized polypeptide drug comprising lyophilizing a solution of the drug in a solvent and milling the lyophilized drug in a fluid energy mill using a pressurized gas, the improvement wherein the polypeptide drug is labile under the milling conditions and the lyophilized drug contains a milling stabilizer.

15

12. The process of claim 11 wherein the milling stabilizer comprises sorbitol, mannitol, sucrose, lactose, trehalose, serum albumin, polyvinylpyrrolidone, or mixtures thereof.

20

13. The process of claim 11 wherein the milling stabilizer is added to the solution.

25

14. The process of claim 13 wherein the concentration of milling stabilizer in the solution is 0.5 to 50 mg/ml.

30

35

INTERNATIONAL SEARCH REPORT

PCT/US 93/00373

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 A61K9/14; A61K9/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP,A,0 384 752 (ELI LILLY AND COMPANY) 29 August 1990	1,4,5,7
Y	see claims 1-4 see page 2, line 3 - line 6 see page 2, line 36 - line 39 see page 3, line 13 - line 45 ---	2,3,8, 10-14
Y	WO,A,9 011 091 (CENTOCOR INC.) 4 October 1990 see claims 1,12 see page 2, line 14 - line 24 see page 4, line 27 - page 5, line 1 -----	2,3,8, 10-14
<p>¹⁰ Special categories of cited documents : ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
18 MARCH 1993	37. 04. 93	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	VENTURA AMAT A.	

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9300373
SA 69620

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18/03/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0384752	29-08-90	US-A- 5021554	04-06-91
		AU-B- 625298	09-07-92
		AU-A- 5006990	30-08-90
		CA-A- 2010328	24-08-90

WO-A-9011091	04-10-90	CA-A- 2049342	28-09-90
		EP-A- 0465513	15-01-92
		JP-T- 4504253	30-07-92
